


# Migratory connectivity in the ocean

Mitigating threats from offshore activities to migratory marine species



Humpback whales in Mo'orea, French Polynesia. Source: Toby Matthews, Ocean Image Bank.

## Key Messages

- Migratory marine species play a pivotal role in the health of the ocean and the ability for people to receive benefits from marine ecosystems.
- Offshore energy production and resource extraction can pose a threat to migratory marine species through noise pollution, oil spills and leaks, collisions and habitat loss, degradation and fragmentation.
- Threats can be mitigated through application of the avoidance and minimisation steps of the mitigation hierarchy, as well as robust environmental impact assessments that take direct and indirect impacts into account.
- Threats to migratory marine species differ through the project lifecycle and so different mitigation measures are needed at different stages.
- Taking part in strategic environmental assessments is a way companies can mitigate cumulative impacts on migratory marine species.
- Many datasets exist that are related to migratory marine species and that companies can use to inform environmental impact assessments, including those containing raw tracking data, areas of importance and key habitat distribution.

## Introduction

Migratory marine species include some of the last remaining megafauna of the ocean. These species help maintain healthy ecosystems and enable people to benefit from ecosystem services, such as fisheries and eco-tourism.

Migratory marine species are highly mobile species that connect disparate areas of the ocean through their movements. As such, they are a form of ecological connectivity. Ecological connectivity describes the “unimpeded movement of species and the flow of natural processes that sustain life on Earth”<sup>1</sup>.

Migratory marine species embark on migrations to make use of resources and conditions that differ in time and space. Migrations vary in duration and distance between different species, with some taking multiple years and covering entire ocean basins. Migratory marine species

include, *inter alia*, whales, sharks, tuna, turtles, and seabirds.

However, migratory marine species are threatened and some currently face extinction. They face a myriad of threats, including from offshore resource extraction and energy production. These activities can adversely impact migratory marine species and jeopardise their continued survival. Yet, these activities are needed to provide the world with a secure, affordable energy supply and natural resources for technology. Companies with offshore activities face the challenge of contributing towards a sustainable society whilst avoiding major impacts on marine biodiversity. This technical briefing explores how this can be achieved, considering offshore resource extraction and energy production.

## Migratory marine species benefit the ocean and people

Migratory marine species often have outsized ecological roles<sup>2,3</sup>. Many are apex predators, meaning they play a vital role in maintaining healthy and productive ecosystems and maintain balance in food webs<sup>4</sup>. As such, many migratory marine species are considered keystone species and their presence, absence or abundance can be used to estimate the occurrence of less detectable species or to infer ecosystem health<sup>5</sup>. For example, presence of sharks on coral reefs can be used to infer the health of the reef<sup>6</sup>.

Migratory marine species also contribute towards nutrient cycling as part of their migrations. Seabirds feed in the open ocean and transport nutrients to nesting sites on islands where they breed. The nutrients leach into the surrounding waters, enhancing coral reef productivity and functioning<sup>7</sup>. Similarly, sea turtle eggs within nests on beaches introduce nutrients

to beach ecosystems<sup>8</sup>, and sharks transfer nutrients from offshore waters to near-shore reefs<sup>9</sup>.

Many migratory marine species also have great cultural and spiritual significance, particularly for indigenous peoples and local communities. Sea turtles have been used for ritual and subsistence purposes for millennia. For example, in Rapa Nui (Easter Island), they are considered protector beings and guides for sailors<sup>10</sup>.

Migratory marine species also benefit local and national economies through eco-tourism. Shark eco-tourism is valued at \$314 million globally per year<sup>11</sup>. Over the course of its lifetime, a whale is estimated to generate \$2 million in economic benefits from eco-tourism, fisheries enhancement, and carbon capture<sup>12</sup>.

# Migratory marine species are threatened with extinction

Migratory marine species include some of the most threatened animals on Earth: 21% are classified as Threatened with extinction and a further 11% are classified as Near Threatened on the IUCN Red List of Threatened Species<sup>13</sup>. This conservation status is caused by a number of pressures, including fishing, shipping, offshore energy production and mining, pollution, habitat loss and degradation, and climate change<sup>13</sup>.

Such pressures are not exclusive to migratory marine species and are drivers of biodiversity loss more broadly. However, these species have characteristics that make them particularly vulnerable. Their migration routes are vast, sometimes covering thousands of kilometres. This exposes them to multiple anthropogenic stressors and can lead to cumulative impacts<sup>14</sup>. In addition, most are large-bodied animals, meaning they are long living and slow to sexually mature. Bowhead whales (*Balaena mysticetus*) can live for up to 200 years<sup>15</sup> and reach maturity

at approximately 20 years of age<sup>16</sup>. This means if a population declines it will take many years to recover. For example, the once-depleted leatherback turtle population in the Atlantic Ocean has shown a gradual recovery after decades of intrusive conservation efforts, yet recovery has taken 2-3 decades<sup>17 18 19</sup>.

The wide-ranging migrations of migratory marine species can make conservation efforts challenging. These species move between Exclusive Economic Zones (EEZs) and the High Seas as they complete their migrations. Efforts to protect them in one area can be undone by a lack of protection elsewhere. Coordinated management along the entirety of migration routes is needed but can be challenging due to the large areas and number of jurisdictions involved<sup>20</sup>. For example, leatherback turtles in the Pacific travel through 32 EEZs as well as the High Seas<sup>21</sup>.

## Offshore activities pose a potential threat

The ocean is highly important for energy production, resource extraction and transportation of goods. It is the world's largest generator of both renewable and non-renewable energy<sup>22</sup>. The ocean's role in energy production is becoming more prominent as the world aims to rapidly transition to renewable energy. Resource extraction through coastal mining provides minerals such as sand to the construction industry. It is used to build roads and buildings, and to make glass and electronics. After water, sand is the second most highly exploited resource in the world<sup>23</sup>. The ocean is also vital for global trade: 80% of world trade is delivered by ships<sup>24</sup>.

However, these offshore activities pose a threat to migratory marine species and can contribute towards population declines if those threats are not managed. Although other anthropogenic activities in the ocean represent more acute threats to certain species groups (e.g., fishing is a particularly large threat to seabirds and turtles), cumulative impacts on migratory marine species from multiple stressors means negative impacts from offshore activities can be more severe than if they occurred in isolation. Table 1 provides an overview of the types of threats from offshore activities, including some emerging threats from novel activities.

Table 1. An overview of threats to migratory marine species from offshore activities.

Threat	Description	<u>Example of direct impact</u>	<u>Example of indirect impact</u>
Habitat loss, degradation fragmentation	<p>Habitat loss, degradation fragmentation can occur from immediate physical alteration of a site for any offshore activity. This can have a pronounced impact if the site is particularly important for a species e.g., a feeding or breeding site. Many shark species depend on coastal or near-shore habitats for their early-life development, such as saltmarshes<sup>25</sup> or mangroves<sup>26</sup>. As such, loss, degradation or fragmentation can affect the availability or effectiveness of habitats that juveniles depend upon<sup>27</sup>. Habitat loss, degradation and fragmentation can result from offshore infrastructure construction, but also from coastal infrastructure which supports offshore activities, such as the construction of ports.</p>	<p>Offshore activities destroy, degrade or fragment a site used by migratory marine species to feed, displacing them from that area.</p>	<p>The fitness of the displaced population decreases as other feeding sites do not have the same abundance of food, or there are a lack of other feeding sites altogether.</p>
Noise pollution and seismic activity	<p>Noise pollution and seismic activity is most often a threat that has been associated with marine mammals<sup>28 29 30</sup>, but it can affect other species including fish<sup>31</sup> and turtles too<sup>32</sup>. Noise pollution is created by, <i>inter alia</i>, ships, mining activities, offshore energy infrastructure construction. Seismic activity stems from activities such as offshore exploration for oil and gas.</p> <p>Noise pollution from anthropogenic activities can 'drown out' natural noises in the ocean or disturb species that are sensitive to sound. This is problematic for those that use sound to navigate or find mates and food. For instance, orcas have been found to spend less time feeding in the presence of boat noise compared to when there is none<sup>33 34</sup>. In a laboratory experiment, fish that shoal were found to be less cohesive and coordinated in their movements when played noises replicating marine construction, which in the wild could make them more vulnerable to predators<sup>35</sup>. Background anthropogenic noise has also been shown to stress whale species, in turn reducing their fitness<sup>36</sup>.</p> <p>Seismic activity used for surveying similarly affects species sensitive to sound. It differs from other noise pollution because it produces high</p>	<p>Noise pollution prevents whale species from hearing mates, increases their stress and causes them to feed less.</p>	<p>The fitness of individuals is reduced and/or they can't locate a mate, meaning they have less offspring and the population declines.</p>

Threat	Description	Example of direct impact	Example of indirect impact
	<p>intensity, high decibel sounds in pulses that can directly injure or kill individuals<sup>37 38</sup>. Seismic activity can alter behaviour, such as causing individuals to dive to deeper waters due to fear<sup>39 40</sup>. It can also affect the ability of marine mammals to navigate and cause large-scale strandings<sup>41</sup>. In fish, seismic activity can rupture their swim bladders, used to control buoyancy<sup>42</sup>.</p>		
Collisions	<p>Some migratory marine species are vulnerable to collisions with particular types of infrastructure used for offshore activities and shipping, causing direct mortality or serious injury. For example, offshore wind turbines are predicted to be a risk to seabirds<sup>43</sup>. Large-bodied species of whale and shark, as well as turtles and seabirds, are vulnerable to ship strikes<sup>44 45 46</sup>.</p>	<p>Ship strikes within an area reduce the local populations of seabirds, turtles and marine mammals.</p>	<p>Eco-tourist numbers in the area decline as sightings of migratory marine species become less frequent.</p>
Oil spills and leaks	<p>Oil spills and leaks degrade water and have an immediate impact by smothering individuals, causing death or seriously affecting health. Offshore oil production is most associated with oil spills, but ships can also release pollutants into the water. Heavy metals can have long-lasting impacts by accumulating in migratory marine species because they are often apex predators or high up in the food chain. Persistent organic compound accumulation is virtually impossible to solve for individuals or pods that already have high levels<sup>47</sup>. For example, some pods of orcas have been unable to produce young in many years due to the high levels of persistent organic compounds they have accumulated in their bodies<sup>48</sup>. The Stockholm Convention on Persistent Organic Pollutants<sup>49</sup> aims to address the issues associated with persistent organic pollutants, yet this threat to marine species, and in particular marine mammals, continues<sup>50</sup>.</p>	<p>Oil spills and leaks smother migratory marine species.</p>	<p>Persistent organic pollutants accumulate in species such as marine mammals, causing infertility and the population to decline as no offspring add to the population.</p>
Electromagnetic fields	<p>Electromagnetic fields are a relatively under-studied threat which is caused by submarine power cables that connect to offshore wind farms producing an electromagnetic field. Changes to migratory fish behaviour has been observed, although it is unknown whether these are temporary behaviour changes or whether there are any long-lasting impacts<sup>51</sup>. Some migratory marine species, such as turtles, are sensitive to the Earth's magnetic field</p>	<p>Electromagnetic fields from submarine power cables affect the navigation of turtle hatchlings after entering the sea.</p>	<p>The navigational abilities of the turtle hatchlings are affected, meaning rather than swimming out past the continental shelf to avoid predation, they remain near</p>

Threat	Description	<u>Example</u> of direct impact	<u>Example</u> of indirect impact
	and use it for navigation. Although not yet studied, there is a potential for electromagnetic fields produced by submarine cables to impact this navigational ability, particularly for turtle hatchlings. They can distinguish small differences in the Earth's magnetic field and could be more likely to be in close proximity to submarine cables after hatching on beaches and moving into coastal waters <sup>52</sup> .		to the coast and are at risk from predation.



An Oceanic Whitetip Shark in Hawaii. Source: Carlos Diaz, Ocean Imagine Bank.

### Direct and indirect impacts

includes examples of both direct and indirect impacts from different threats posed by offshore activities on migratory marine species. Whereas direct impacts are more obvious and easier to study, indirect impacts are more nuanced and harder to monitor<sup>53</sup>. Direct impacts on migratory marine species can lead to indirect impacts on the wider ecosystem<sup>54 55 56</sup> and people<sup>57 58 59</sup>, in addition to migratory marine species themselves.

Indirect impacts include displacement as key habitats are removed, degraded, or fragmented, or as species seek to avoid sites due to proximity of anthropogenic activities<sup>37 60</sup>. This means they are not able to access key sites for feeding or breeding and expend extra energy taking longer routes to other sites<sup>61</sup>. In turn, this can result in

reduced health and fitness amongst individuals and populations, making them more susceptible to predators and disease. Offshore activities can also cause prey disturbance, displacing prey that migratory marine species rely on.

### Potential positive impacts

In some cases, offshore activities can result in positive impacts for migratory marine species too. Certain offshore infrastructure can support new feeding sites for migratory marine species through creating new habitats which marine species have colonised, such as wind farms acting as artificial reefs<sup>62 63</sup>. In addition, offshore sites such as wind farms can offer shelter for migratory marine species (amongst other marine life) from threats such as fishing or shipping<sup>62</sup>.

## Threats from offshore activities can be avoided and minimised

The mitigation hierarchy can be used to mitigate threats to migratory marine species from offshore activities (Table 2). It is particularly important to prioritise the **avoidance** and **minimisation** steps, because these are typically vulnerable species.

**Restoration** of populations of migratory marine species is highly challenging due to their longevity<sup>15 64</sup>. For example, certain whale populations are only now just recovering from hunting in the 19<sup>th</sup> and 20<sup>th</sup> centuries<sup>65</sup>, highlighting the time needed for population recovery of these long-lived, slow-to-mature species. These features are common characteristics of many migratory marine species<sup>16 64</sup>. Due to these challenges, restoration of marine habitats that migratory marine species depend on, and that have been lost, degraded or fragmented as a result of offshore activities, can be used as a proxy means of migratory marine species restoration. This may be possible in

some cases, although not all. Restoration may not be appropriate if the site has an outsized role in supporting migratory marine species and its removal would jeopardise their continued survival. Table 2 outlines how restoration could be undertaken in the context of migratory marine species populations and the habitats they rely on.

**Offsetting** impacts on migratory marine species populations is rarely likely to be legitimate or effective due to the inherent challenges of restoring migratory marine species populations, as outlined under the restoration step<sup>66</sup>. Offsetting impacts to marine habitats as a proxy may be possible if there was strong evidence migratory marine species would benefit and if the offsetting met key criteria (outlined in Table 2). However, offsetting is still highly challenging in the context of migratory marine species and further research is needed as to how to ensure it would be legitimate in the context of these species.

Table 2. Examples of how the mitigation hierarchy can be applied in the context of migratory marine species.

Step of mitigation hierarchy	Application of step in the context of different threats, using examples.
Avoidance	<p><u>Threat:</u> Placement of offshore activities on key sites e.g., feeding, breeding or nursery grounds (<i>habitat loss, degradation and fragmentation</i>).</p> <p><u>Action:</u> <b>Avoid</b> placing offshore activities over those key sites that are essential to the continued survival of a species.</p>
Minimisation	<p><u>Threat:</u> Seismic surveys for oil and gas during the exploration stage (<i>noise pollution and seismic surveys</i>).</p> <p><u>Action:</u> Halt surveys during the portion of the year when migratory marine species are using/passing through the area to <b>minimise</b> impact (e.g., when turtles are gathering offshore to breed and nest).</p> <p><u>Threat:</u> Shipping to support offshore energy production and resource extraction (<i>collisions</i>).</p> <p><u>Action:</u> Re-route shipping lanes to <b>minimise</b> overlap with known migratory pathways and/or where there is overlapping, reducing ship speeds.</p>
Restoration	<p><u>Threat:</u> Oil spills and leaks from offshore oil extraction (<i>oil spills and leaks</i>).</p> <p><u>Action:</u> (Restoring migratory marine species populations) Provide additional protection to local seabird nesting sites to <b>increase/restore</b> the population. (Restoring habitats migratory marine species rely on) Oil spill clean-up and active <b>restoration</b> of degraded habitats migratory marine species use e.g., planting seagrass seeds.</p>
Offsetting	<p>Jacob et al. (2020)<sup>67</sup> outline the following criteria that marine offsets need to meet in order to be legitimate:</p> <ul style="list-style-type: none"> <li>• Comparable (a true exchange for the biodiversity lost)</li> <li>• Additional (over and above what would have happened without the offset)</li> <li>• Lasting (in place for at least as long as the impacts)</li> <li>• Within limits (some impacts are not offsetable)</li> </ul> <p>Further research is needed to establish what offsets mean in the context of migratory marine species and if they are possible.</p>

The mitigation hierarchy can be applied as part of a robust environmental impact assessment (EIA)<sup>68</sup>. It should be applied iteratively and at each stage of the operation or project lifecycle, including exploration, development, operation, and decommissioning<sup>69</sup>. Different stages of an operation or project lifecycle can pose different threats to migratory marine species. This means different steps of the mitigation hierarchy may need to be applied at each stage and different actions are needed to mitigate threats. For example, noise pollution is a threat from offshore

oil and gas during the exploratory phase when seismic surveys are carried out. During the operational phase, oil spills and leaks pose a larger threat.

EIAs and biodiversity action plans should reflect the dynamic nature of migratory marine species in space and time. In addition to different actions to mitigate threats being needed at different stages of the project lifecycle, different actions may be needed within the same stage and in different months or seasons to account for

migratory marine species presence or absence during different parts of the year.

Because migratory marine species are a form of connectivity, it means negative impacts on their populations can be far-reaching and felt in disparate locations to where the stressor occurs<sup>70</sup>. EIAs should take this into account when defining the area of influence, as much as they are able to. For example, if a whale shark population declines on one side of an ocean, those individuals never complete their onward journey to other locations, where they are not present to feed on plankton. The absence of whale sharks could result in increased plankton blooms, reducing productivity of fisheries in the area. Although incorporating such indirect impacts into an EIA is challenging, it is a step towards ensuring all potential threats are truly mitigated.

To address the issue of cumulative impacts on migratory marine species from multiple sectors,

companies that operate offshore activities could become involved in a strategic environmental assessment (SEA). They assess the potential effects of all proposed activities within a given area at the planning stage and are often used in marine spatial planning processes. SEAs can use cumulative impact assessments to calculate cumulative impacts within a given area<sup>71</sup>. Involvement in a SEA process could help companies that operate offshore understand their impact on migratory marine species in relation to other ocean activities. In addition, the information provided would be specific to the geographic area and its context, rather than having to rely on generalisations.

Companies may also find that certain actions, such as participating in a SEA, become an emerging requirement in the coming decade in light of the adoption of the Kunming-Montreal Global Biodiversity Framework<sup>72</sup> under the Convention on Global Biodiversity.

## Data and knowledge

Mitigating the threats to migratory marine species from offshore activities requires data and knowledge. Tracking data has increased over the past two decades<sup>73</sup>, and more recently, so have tools which synthesise this data to inform decision making.

Migratory marine species datasets that consist of raw tracking data include the [Seabird Tracking Database](#), which contains the world's largest collection of seabird tracking data. Others include the [Global distribution of Sei whales](#) and the [Global distribution of Sperm whales](#). Some datasets are based on important feeding or breeding sites, such as the [Global Distribution of Sea Turtle Nesting Sites](#). The [Migratory Connectivity in the Ocean \(MiCO\)](#)<sup>74</sup> system aggregates tracking data of migratory marine species, so can be used to consider multiple species. Datasets such as these can be used to determine whether offshore activities overlap with migratory marine species.

Other datasets can be used that identify priority sites for migratory marine species. The [Important Marine Mammal Area \(IMMA\)](#) database identifies areas of importance to the conservation of marine mammal species, including migratory species. A similar [Important Marine Turtle Area](#) database is currently being developed to identify areas of particular significance for the conservation of marine turtles and their contributions to local people, including habitats used for migration. The [Key Biodiversity Areas \(KBA\)](#) dataset can be filtered by the species that triggered the KBA, enabling filtering of KBAs triggered by migratory marine species.

Other tools of relevance to offshore energy production include the [Avian Sensitivity Tool for Energy Planning \(AVISTEP\)](#), which shows the sensitivity of birds to energy infrastructure and includes offshore wind as well as seabirds.

Other non-migratory marine species spatial datasets can be used in EIAs to help manage impacts on these species, such as spatial data on

the habitats they rely on. [Ocean+ Habitats](#) and [Ocean+ Data Viewer](#) contain data on the distribution of critical marine and coastal habitats, and broader marine biodiversity spatial datasets, respectively. [UN Biodiversity Lab](#) contains global biodiversity spatial data, including marine-specific datasets.

As well global datasets, local, national or regional datasets of both species and habitats are important to include in EIAs, where available. They will likely provide higher levels of detail for specific sites, complementing global datasets which reflect the vast ranges migratory species encompass.

Knowledge garnered by indigenous peoples (IPs) and local communities (LCs) is also a useful and powerful resource. The cultural significance of migratory marine species to IPs and LCs mean

they have a wealth of knowledge and understanding of these species that has been built up over many generations. As well as complementing existing datasets, IPs and LCs knowledge can be used in areas where there is limited or gaps in data. Further still, IPs and LCs knowledge can be used as a source of the values of migratory marine species and as such, a measure of the impacts of offshore activities that need to be taken into account within EIAs. Data and knowledge on migratory marine species can be used to inform project screening and EIAs when projects are in the planning stage. However, they can also be applied to operational projects during ongoing or future phases of the project lifecycle. Field surveys, as well as consultation with migratory marine species and regional experts, can be used to complement datasets and inform decision-making.

## Policy relevance

Companies will often be operating in countries that are a Party to international conventions aiming to conserve and sustainably use biodiversity, including the Convention on Biological Diversity<sup>75</sup> and the Convention on the International Trade in Endangered Species of Wild Fauna and Flora<sup>76</sup>. Of particular pertinence to migratory marine species is the Convention on the Conservation of Migratory Species of Wild Animals (CMS)<sup>77</sup>. Parties to the CMS have agreed to conserve and restore habitats that migratory species rely on, as well as to “prevent, remove, compensate for or minimize...activities or obstacles that seriously impede or prevent the migration of...species”<sup>78</sup>. By operating within a country that is Party to the CMS, companies have an obligation and important role to ensure this commitment is met.

The CMS operates as an umbrella agreement, under which specific taxonomic and/or regional

agreements and memoranda of understanding exist. Those relating to migratory marine species include, *inter alia*, the Agreement on the Conservation of Cetaceans of the Black Sea, Mediterranean Sea and contiguous Atlantic Area<sup>79</sup> and the Agreement on the Conservation of Small Cetaceans of the Baltic, North East Atlantic, Irish and North Seas<sup>80</sup>. In addition, single species action plans exist under the CMS as well as its daughter agreements, such the Conservation Plan for Migratory Sharks<sup>81</sup> and the Single Species Action Plan for the Angel Shark in the Mediterranean Sea<sup>82</sup>.

The CMS produces guidelines to support its Parties, some of which may be useful for companies wishing to mitigate their impacts on migratory marine species. For example, guidelines on marine noise-generating activities within environmental impact assessments<sup>83</sup>.

## Conclusion

Migratory marine species are keystones of ocean health. They help maintain healthy, balanced ecosystems and, in turn, the ability for people to benefit from the ocean. But they are also highly threatened. Whilst their migratory behaviour allows them to access resources that change in time and space, it also exposes them to numerous stressors that has been described as 'death by a thousand cuts'.

All ocean-users have a role to play in mitigating these threats. There are actions that organisations with offshore activities can take to mitigate potential threats. Data to inform these mitigation actions is also increasing, including rapid

increases in availability of species tracking data, and these can be used to inform measures to mitigate harm.

With the adoption of the Kunming-Montreal Global Biodiversity Framework (GBF), countries will be working to revise and update their national biodiversity strategies and action plans and take action to meet the Kunming-Montreal 2030 Global Targets. Efforts by the private sector to protect biodiversity will be key to the GBF's success. With migratory marine species including some of the most threatened species of Earth, efforts to protect them can provide a fundamental contribution towards the success of the GBF.

**Citation:** UNEP-WCMC (2022). Migratory connectivity in the ocean. Cambridge, UK.

**Authors:** Holly Brooks and Nina Bhola

**Acknowledgements:** The authors would like to thank reviewers including Cleo Cunningham, Stacey Baggaley and Giulia Costa-Domingo

**Available online at:** [www.proteuspartners.org](http://www.proteuspartners.org)

**Contact:** Stacey Baggaley ([stacey.baggaley@unep-wcmc.org](mailto:stacey.baggaley@unep-wcmc.org))

Copyright: 2023 United Nations Environment Programme

The UN Environment Programme World Conservation Monitoring Centre (UNEP-WCMC) is a global Centre of excellence on biodiversity. The Centre operates as a collaboration between the UN Environment Programme and the UK-registered charity WCMC. Together we are confronting the global crisis facing nature.

**Copyright release:** This publication may be reproduced for educational or non-profit purposes without special permission, provided acknowledgement to the source is made. Reuse of any figures is subject to permission from the original rights holders. No use of this publication may be made for resale without permission in writing from UN Environment Programme. Applications for permission, with a statement of purpose and extent of reproduction, should be sent to the Director, UNEP-WCMC, 219 Huntingdon Road, Cambridge, CB3 0DL, UK.

**Disclaimer:** The contents of this report do not necessarily reflect the views or policies of the UN Environment Programme, contributory organisations or editors. The designations employed and the presentations of material in this report do not imply the expression of any opinion whatsoever on the part of the UN Environment Programme or contributory organisations, editors or publishers concerning the legal status of any country, territory, city area or its authorities, or concerning the delimitation of its frontiers or boundaries or the designation of its name, frontiers or boundaries. The mention of a commercial entity or product in this publication does not imply endorsement by the UN Environment Programme.

<http://www.unep-wcmc.org/>

## References

- <sup>1</sup> Convention on Migratory Species (2020b). Improving ways of addressing connectivity in conservation of migratory species. COP Resolution 12.26 (Rev COP13) Available from: <https://cutt.ly/4x2AU0S>
- <sup>2</sup> Bornatowski, H., Navia, A. F., Rennó Braga, R., Abilhoa, V., and Maia Correa, M. F. (2014). Ecological importance of sharks and rays in a structural foodweb analysis in southern Brazil. *ICES J. Mar. Sci.* 71, 1586–1592. doi: 10.1093/icesjms/fsu025
- <sup>3</sup> Roman, J., Estes, J. A., Morissette, L., Smith, C., Costa, D., McCarthy, J., et al. (2014). Whales as marine ecosystem engineers. *Front. Ecol. Environ.* 12, 377–385. doi: 10.1890/130220
- <sup>4</sup> Schefer, M., Carpenter, S. and de Young, B. (2005). Cascading effects of overfishing marine systems. *Trends in Ecology & Evolution*, 20 (11): 579-581
- <sup>5</sup> Hazen, E.L., Abrahms, B., Brodie, S., Carroll, G., Jacox, M.G., Savoca, M.S., Scales, K.L., Sydeman, W.J. and Bograd, S.J. (2019). Marine top predators as climate and ecosystem sentinels. *Frontiers in Ecology and the Environment*, 17 (10): 565-574
- <sup>6</sup> Mukharror, D.A., Tiara Baiti, I., Ichsan, M., Pridina, N. and Triutami, S. (2017). Correlation between existence of reef sharks with abundance of reef fishes in south waters of Morotai Island (North Moluccas). *IOP Conf. Ser.: Earth Environ. Sci.*, 89: DOI 10.1088/1755-1315/89/1/012006
- <sup>7</sup> Graham, N.A.J., Wilson, S.K., Carr, P., Hoey, A.S., Jennings, S. and MacNeil, M.A. (2018) Seabirds enhance coral reef productivity and functioning in the absence of invasive rats. *Nature*, 559:250–253
- <sup>8</sup> Lovich, J.E., Enne, J.R., Agha, M. and Gibbons, J.W. (2018). Where have all the turtles gone, and why does it matter? *BioScience*, 68 (10): 771-781
- <sup>9</sup> Williams, J.J., Papastamatiou, Y.P., Caselle, J.E., Bradley, D. and Jacoby, D.M.P. (2018). Mobile marine predators: an understudied source of nutrients to coral reefs in an unfished atoll. *Proceedings of the Royal Society B*, 285 (1875): <https://doi.org/10.1098/rspb.2017.2456>
- <sup>10</sup> Alvarez-Varas, R., Barrios-Garrido, H., Skamiotis-Gomez, I. and Petitpas, R. (2020). Cultural role of sea turtles on Rapa Nui (Easter Island): Spatial and temporal contrast in the Pacific island region. *Island Studies Journal*, 15 (1): 253-270
- <sup>11</sup> Cisneros-Montemayor, A.M., Barnes-Mauthe, M., Al-Abdulrazzak, D., Navarro-Holm, E. and Sumaila, U.R. (2013). Global economic value of shark ecotourism: implications for conservation. *Oryx*, 47 (3): 381-388
- <sup>12</sup> Chami, R., Cosimano, T., Fullenkamp, C., Berzaghi, F., Espanol-Jimenez, S., Marcondes, M. and Palazzo, J. (2022). The value of nature to our health and economic well-being: A framework with applications to elephants and whales. In: Paganetto, L. (eds) *Economic Challenges for Europe After the Pandemic*. Springer Proceedings in Business and Economics. Springer, Cham. [https://doi.org/10.1007/978-3-031-10302-5\\_7](https://doi.org/10.1007/978-3-031-10302-5_7)
- <sup>13</sup> Lascelles, B., Di Sciara, G.N., Agardy, T., Cuttelod, A., Eckert, S., Glowka, L., Hoyt, E., Llewellyn, F., Louzao, M., Ridoux, V. and Tetley, M. (2014). Migratory marine species: their status, threats and conservation management needs. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24 (2): 111-127
- <sup>14</sup> Maxwell, S.M., Hazen, E.L., Bograd, S.J., Halpern, B.S., Breed, G.A., Nickel, B., Teutschel, N.M., Crowder, L.B., Benson, S., Dutton, P.H., Bailey, H., Kappes, M.A., Kuhn, C.E., Weise, M.J., Mate, B., Shaffer, S.A., Hassrick, J.L., Henry, R.W., Irvine, L., McDonald, B.I., Robinson, P.W., Block, B.A. and Costa, D.P. (2013). Cumulative human impacts on marine predators. *Nature Communications*, 4 (2699)
- <sup>15</sup> Seim, I., Ma, S., Zhou, X., Gerashchenko, M.V., Lee, S-G., Suydam, R., George, J.C., Bickham, J.W. and Gladyshev, V.N. (2014). The transcriptome of the bowhead whale *Balaena mysticetus* reveals adaptations of the longest-lived mammal. *Aging (Albany NY)*, 6 (10): 879-899
- <sup>16</sup> Schell, D.M., Saupe, S.M. and Haubenstock, N. (1989). Bowhead whale (*Balaena mysticetus*) growth and feeding as estimated by  $\delta^{13}C$  techniques. *Marine Biology*, 103: 433-443
- <sup>17</sup> Dutton, D.L., Dutton, P.H., Chaloupka, M. and Boulon, R.H. (2005). Increase of a Caribbean leatherback turtle *Dermochelys coriacea* nesting population linked to long-term nest protection. *Biological Conservation*, 126 (2): 186-194
- <sup>18</sup> Stewart, K.R., Martin, K.J., Johnson, C., Desjardin, N., Eckert, S.A. and Crowder, L.B. (2014). Increased nesting, good survival and variable site fidelity for leatherback turtles in Florida, USA. *Biological Conservation*, 176: 117-125
- <sup>19</sup> Stewart, K., Sims, M., Meylan, A., Witherington, B., Brost, B. and Crowder, L.B. (2011). Leatherback nests increasing significantly in Florida, USA; trends assessed over 30 years using multilevel modelling. *Ecological Applications*, 21 (1): 263-273
- <sup>20</sup> Dunn, D.C., Harrison, A-L., Curtice, C., DeLand, S., Donnelly, B., Fujioka, E., Heywood, E., Kot, C.Y., Poulin, S., Whitten, M., Akesson, S., Alberini, A., Appeltans, W., Arcos, J.M., Bailey, H., Ballance, L.T., Block, B., Blondin, H., Boustany, A.M., Brenner, J., Catry, P., Cejudo, D., Cleary, J., Corkeron, P., Costa, D.P., Coyne, M., Ortuno Crespo, G., Davies, T.E., Dias, M.P., Douvere, F., Ferretti, F., Formia, A., Freestone, D., Friedlaender, A.S., Frisch-Nwakanma, H., Barrio Frojan, C., Gjerde, K.M., Glowka, L., Godley, B.J., Gonzalez-Solis, J., Pedro Granadeiro, J., Gunn, V., Hashimoto, Y., Hawkes, L.M., Hays, G.C., Hazin, C., Jimenez, J., Johnson, D.E., Luschi, P., Maxwell, S.M., McClellan, C., Modest, M., Notarbartolo di Sciara, G., Herrero Palacio, A., Palacios, D.M., Pauly, A.,

- Rayner, M., Rees, A.F., Ross Salazar, E., Secor, D., Sequeira, A.M.M., Spalding, M., Spina, F., Van Parijs, S., Wallace, B., Varo-Cruz, N., Virtue, M., Weimerskirch, H., Wilson, L., Woodward, B. and Halpin, P.N. (2019). The importance of migratory connectivity for global ocean policy. *Proc. R. Soc. B*, 286: 20191472
- <sup>21</sup> Harrison, A.-L., Costa, D.P., Winship, A.J., Benson, S.R., Bograd, S.J., Antolos, M., Carlisle, A.B., Dewar, H., Hutton, P.H., Jorgensen, S.J., Kohin, S., Mate, B.R., Robinson, P.W., Schaefer, K.M., Shaffer, S.A., Shillinger, G.L., Simmons, S.E., Weng, K.C., Gjerde, K.M. and Block, B.A. (2018). The political biogeography of migratory marine predators. *Nature Ecology & Evolution*, 2: 1571-1578
- <sup>22</sup> World Economic Forum (2022). What ocean sustainability means for business. White paper, November 2022
- <sup>23</sup> UNEP (2022). Sand and sustainability: 10 strategic recommendations to avert a crisis. GRID-Geneva, United Nations Environment Programme, Geneva, Switzerland
- <sup>24</sup> UNCTAD (2022). Review of Maritime Transport 2022. UNCTAD/RMT/2022
- <sup>25</sup> Niella, Y., Raoult, V., Gaston, T., Goodman, K., Harcourt, R., Peddemors, V. and Smoothey, A.F. (2022). Reliance of young sharks on threatened estuarine habitats for nutrition implies susceptibility to climate change. *Estuarine, Coastal and Shelf Science*, 268: 107790
- <sup>26</sup> White, W.T. and Potter, I.C. (2004). Habitat partitioning among four elasmobranch species in nearshore, shallow waters of a subtropical embayment in Western Australia. *Marine Biology*, 145: 1023-1032
- <sup>27</sup> Sievers, M., Brown, C.J., Tulloch, V.J.D., Pearson, R.M., Haig, J.A., Turschwell, M.P. and Connolly, R.M. (2019). The role of vegetated coastal wetlands for marine megafauna conservation. *Trends in Ecology & Evolution*, 34 (9): 807-817
- <sup>28</sup> Caldwell, J. (2004). Are seismic air-gun sources harmful to marine mammals? *Polarforschung*, 72: 103-107
- <sup>29</sup> Gordon, J., Gillespie, D., Potter, J., Simmonds, M.P. and Thompson, D. (2003). A review of the effects of seismic surveys on marine mammals. *Marine Technology Society Journal*, 37 (4): 16-34
- <sup>30</sup> Weilgart, L. (2007). A brief review of known effects of noise on marine mammals. *International Journal of Comparative Psychology*, 20 (2)
- <sup>31</sup> Di Franco, E., Pierson, P., Di Lorio, L., Calo, A., Cottalorda, J.M., Derijard, B., Di Franco, A., Galve, A., Guibbolini, M., Lebrun, J., Micheli, F., Priouzeau, F., Risso-de Faverney, C., Rossi, F., Sabourault, C., Spennato, G., Varrando, P. and Guidetti, P. (2020). Effects of marine noise pollution on Mediterranean fishes and invertebrates: A review. *Marine Pollution Bulletin*, 159
- <sup>32</sup> Nelms, S.E., Piniak, W.E.D., Weir, C.R. and Godley, B.J. (2016). Seismic surveys and marine turtles: An underestimated global threat? *Biological Conservation*, 193: 49-65
- <sup>33</sup> Lusseau, D., Bain, D.E., Williams, R. and Smith, J.C. (2009). Vessel traffic disrupts the foraging behavior of southern resident killer whales *Orcinus orca*. *Endangered Species Research*, 6: 211-221
- <sup>34</sup> Williams, R., Lusseau, D. and Hammond, P.S. (2006). *Biological Conservation*, 113 (3): 301-311
- <sup>35</sup> Herbert-Read, J.E., Kremer, L., Bruintjes, R., Radford, A.N. and Ioannou, C.C. (2017). Anthropogenic noise pollution from pile-driving disrupts the structure and dynamics of fish shoals. *Proceedings of the Royal Society B*, 284 (1863)
- <sup>36</sup> Rolland, R.M., Parks, S.E., Hunt, K.E., Castellote, M., Corkeron, P.J., Nowacek, D.P., Wasser, S.K. and Kraus, S.D. (2012). Evidence that ship noise increases stress in right whales. *Proceedings of the Royal Society B*, 279 (1737)
- <sup>37</sup> Kavanagh, A.S., Nykanen, M., Hunt, W., Richardson, N. and Jessopp, M.J. (2019). Seismic surveys reduce cetacean sightings across a large marine ecosystem. *Scientific Reports*, 9 (19164)
- <sup>38</sup> Schrope, M. (2002). Whale deaths caused by US Navy sonar. *Nature*, 415:106. doi: 10.1038/415106a
- <sup>39</sup> Williams, T.M., Blackwell, S.B., Tervo, O., Garde, E., Sinding, M.-H., S., Richter, B. and Heide-Jorgensen, M.P. (2022). Physiological responses of narwhals to anthropogenic noise: A case study with seismic airguns and vessel traffic in the Arctic. *Functional Ecology*, 36 (9): 2251-2266
- <sup>40</sup> DeRuiter, S.L. and Doukara, K.L. (2012). Loggerhead turtles dive in response to airgun sound exposure. *Endangered Species Research*, 16: 55-63
- <sup>41</sup> Jepson, P.D., Deaville, R., Acevedo-Whitehouse, K., Barnett, J., Brownlow, A., Brownell Jr., R.L., Clare, F.C., Davison, N., Law, R.J., Loveridge, J., Macgregor, S.K., Morris, S., Murphy, S., Penrose, R., Perkins, M.W., Pinn, E., Seibel, H., Siebert, U., Sierra, E., Simpson, V., Tasker, M.L., Tregenza, N., Cunningham, A.A. and Fernandez, A. (2013). What caused the UK's largest common dolphin (*Delphinus delphis*) mass stranding event? *PLoS ONE*, 8 (4): e60953
- <sup>42</sup> Halvorsen, M.B., Casper, B.M., Woodley, C.M., Carlson, T.J. and Popper, A.N. (2012). Threshold for onset of injury in Chinook Salmon from exposure to impulsive pile driving sounds. *PLoS ONE*, 7 (6): e38968
- <sup>43</sup> Furness, R.W., Wade, H.M. and Masden, E.A. (2013). Assessing vulnerability of marine bird populations to offshore wind farms. *Journal of Environmental Management*, 119: 56-66
- <sup>44</sup> Wormersley, F.C., Humphries, N.E., Queiroz, N., Vedor, M., da Costa, I., Furtado, M., Tyminski, J.P., Abrantes, K., Araujo, G., Bach, S.S., Barnett, A., Berumen, M.L., Bessudo Lion, S., Braun, C.D., Clingham, E., Cochran, J.E.M., de la Parra, R., Diamant, S., Dove, A.D.M., Dudgeon, C.L., Erdmann, M.V., Espinoza, E., Fitzpatrick, R., Gonzalez Cano, J., Green, J.R., Guzman, H.M.,

- Hardenstine, R., Hasan, A., Hazin, F.H.V., Hearn, A.R., Hueter, R.E., Jaidah, M.Y., Labaja, J., Ladino, F., Macena, B.C.L., Morris Jr., J.J., Norman, B.M., Penaherrera-Palma, C., Pierce, S.J., Quintero, L.M., Ramirez-Macias, D., Reynolds, S.D., Richardson, A.J., Robinson, D.P., Rohner, C.A., Rowat, D.R.L., Sheaves, M., Shivji, M.S., Sianipar, A.B., Skomal, G.B., Soler, G., Syakurachman, I., Thorrold, S.R., Webb, D.H., Wetherbee, B.M., White, T.D., Clavelle, T., Kroodsma, D.A., Thums, M., Ferreira, L.C., Meekan, M.G., Arrowsmith, L.M., Lester, E.K., Meyers, M.M., Peel, L.R., Sequeira, A.M.M., Eguiluz, V.M., Duarte, C.M. and Sims, D.W. (2022). Global collision-risk hotspots of marine traffic and the world's largest fish, the whale shark. *PNAS*, 119 (20)
- <sup>45</sup> Hazel, J., Lawler, I.R., Marsh, H and Robson, S. (2007). Vessel speed increases collision risk for the green turtle *Chelonia mydas*. *Endangered Species Research*, 3 (2)
- <sup>46</sup> Coleman, J., Hollyman, P.R., Black, A. and Collins, M.A. (2022). Blinded by the light: Seabird collision events in South Georgia. *Polar Biology*, 45: 1151-1156
- <sup>47</sup> Desforges, J-P., Hall, A., McConnell, B., Rosing-Asvid, A., Barber, J.L., Brownlow, A., De Guise, S., Eulaers, I., Jepson, P.D., Letcher, R.J., Levin, M., Ross, P.S., Samarra, F., Vikingson, G., Sonne, C. and Dietz, R. (2018). Predicting global killer whale population collapse from PCB pollution. *Science*, 361 (6409): 1373-1376
- <sup>48</sup> Jepson, P.D., Deaville, R., Barber, J.L., Aguilar, A., Borrell, A., Murphy, S., Barry, J., Brownlow, A., Barnett, J., Berrow, S., Cunningham, A.A., Davison, N.J., ten Doeschate, M., Esteban, R., Ferreira, M., Foote, A.D., Genov, T., Gimenez, J., Loveridge, J., Llavona, A., Martin, V., Maxwell, D.L., Papachlitzou, A., Penrose, R., Perkins, M.W., Smith, B., de Stephanis, R., Tregenza, N., Verborgh, P., Fernandez, A. and Law, R.L. (2016). PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Scientific Reports*, 6 (18573)
- <sup>49</sup> Stockholm Convention on Persistent Organic Pollutants
- <sup>50</sup> Stuart-Smith, S.J. and Jepson, P.D. (2017). Persistent threats need persistent counteraction: Responding to PCB pollution in marine mammals. *Marine Policy*, 84: 69-75
- <sup>51</sup> Gill, A.B., Bartlett, M. and Thomsen, F. (2012). Potential interactions between diadromous fishes of U.K. conservation importance and the electromagnetic fields and subsea noise from marine renewable energy developments. *Journal of Fish Biology*, 81 (2): 664-695
- <sup>52</sup> Yalcin-Ozdilek, S. and Yalcin, S. (2012). Wind Energy Plants and Possible Effects on Samandag Sea Turtles. *Marine Turtle Newsletter*, 133: 7-9
- <sup>53</sup> UNEP-WCMC (2022). Area of Influence of Site-based operations-Indirect Impacts. Cambridge, UK.
- <sup>54</sup> Atwood, T.B. and Hammill, E. (2018). The importance of marine predators in the provisioning of ecosystem services by coastal plant communities. *Frontiers in Plant Science*, 9
- <sup>55</sup> Heithaus, M., Frid, A., Wirsing, A.J. and Worm, B. (2008). Predicting ecological consequences of marine top predator declines. *Trends in Ecology & Evolution*, 23 (4): 202-210
- <sup>56</sup> Britten, G.L., Dowd, M., Minto, C., Ferretti, F., Boero, F. and Lotze, H.K. (2014). Predator decline leads to decreased stability in a coastal fish community. *Ecology Letters*, 17 (12): 1514-1525
- <sup>57</sup> Alvarez-Varas, R., Barrios-Garrido, H., Skamiotis-Gomez, I. and Petitpas, R. (2020). Cultural role of sea turtles on Rapa Nui (Easter Island): Spatial and temporal contrast in the Pacific island region. *Island Studies Journal*, 15 (1): 253-270
- <sup>58</sup> Vierros, M.K., Harrison, A-L., Sloat, M.R., Ortuno Crespo, G., Moore, J.W., Dunn, D.C., Ota, Y., Cisneros-Montemayor, A.M., Shillinger, G.L., Kehaulani Watson, T. and Govan, H. (2020). Considering Indigenous Peoples and local communities in governance of the global ocean commons. *Marine Policy*, 119
- <sup>59</sup> Reynolds, SD, Norman, BM, Beger, M, Franklin, CE, Dwyer, RG. Movement, distribution and marine reserve use by an endangered migratory giant. *Divers Distrib*. 2017; 23: 1268– 1279. <https://doi.org/10.1111/ddi.12618>
- <sup>60</sup> Brandt, M.J., Diederichs, A., Betke, K. and Nehls, G. (2011). Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series*, 421: 205-216
- <sup>61</sup> Forney, K.A., Southall, B.L., Slooten, E., Dawson, S., Read, A.J., Baird, R.W. and Brownell Jr., R.L. (2017). Nowhere to go: noise impact assessments for marine mammal populations with high site fidelity. *Endangered Species Research*, 32: 391-413
- <sup>62</sup> Lindeboom, H.J., Kouwenhoven, H.J., Bergman, M.J.N., Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., de Hann, D., Dirksen, S., van Hal, R., Hille Ris Lambers, R., ter Hofstede, R., Krijgsveld, K.L., Leopold, M. and Scheidat, M. (2011). Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. *Environmental Research Letters*, 6 (3): 035101
- <sup>63</sup> Hammar, L., Perry, D. and Gullstrom, M. (2016). Offshore wind power for marine conservation. *Open Journal of Marine Science*, 6 (1)
- <sup>64</sup> Mayne, B., Tucker, A.D., Berry, O. and Jarman, S. (2020). Lifespan estimation in marine turtles using genomic promoter CpG density. *PLoS ONE*, 15 (7): e0236888
- <sup>65</sup> Tulloch, V.J.D., Plaganyi, E.E., Matear, R., Brown, C.J. and Richardson, A.J. (2017). Ecosystem modelling to quantify the impact of historical whaling on Southern Hemisphere baleen whales. *Fish and Fisheries*, 19 (1): 117-137

- 
- <sup>66</sup> Vaissiere, A-C., Levrel, H., Pioch, S. and Carlier, A. (2014). Biodiversity offsets for offshore wind farm projects: The current situation in Europe. *Marine Policy*, 48: 172-183
- <sup>67</sup> Jacob, C., van Bochove, J-W., Livingstone, S., White, T., Pilgrim, J. and Bennun, L. (2020). Marine biodiversity offsets: Pragmatic approaches toward better conservation outcomes. *Conservation Letters*, 13 (3)
- <sup>68</sup> The Biodiversity Consultancy (2015). A cross-sector guide for implementing the mitigation hierarchy. Cambridge, UK
- <sup>69</sup> Bennun, L., van Bochove, J., Ng, C., Fletcher, C., Wilson, D., Phair, N., Carbone, G. (2021). Mitigating biodiversity impacts associated with solar and wind energy development. Guidelines for project developers. Gland, Switzerland: IUCN and Cambridge, UK: The Biodiversity Consultancy.
- <sup>70</sup> Jonsson, P.R., Hammar, L., Wåhlström, I., Pålsson, J., Hume, D., Almroth-Rosell, E. and Mattsson, M. (2020). Combining seascape connectivity with cumulative impact assessment in support of ecosystem-based marine spatial planning. *Journal of Applied Ecology*, 58: 576-586
- <sup>71</sup> Jonsson, P.R., Hammar, L., Wåhlström, I., Pålsson, J., Hume, D., Almroth-Rosell, E. and Mattsson, M. (2020). Combining seascape connectivity with cumulative impact assessment in support of ecosystem-based marine spatial planning. *Journal of Applied Ecology*, 58: 576-586
- <sup>72</sup> CBD (2022). CBD/COP/15/L.25
- <sup>73</sup> Hussey, N.E., Kessel, S.T., Aarestrup, K., Cooke, S.J., Cowley, P.D., Fisk, A.T., Harcourt, R.G., Holland, K.N., Iverson, S.J., Kocik, J.F., Mills Fleming, J.E. and Whoriskey, F.G. (2015). Aquatic animal telemetry: A panoramic window into the underwater world. *Science*, 348 (6240): DOI: 10.1126/science.1255642
- <sup>74</sup> Dunn, D.C., Curtice, C., DeLand, S., Donnelly, B., Fujioka, E., Halpin, P., Kot, C. and Nisthar, D. (2020). Advancing understanding of Migratory Connectivity in the Ocean (MiCO). 10pp.
- <sup>75</sup> <https://www.cbd.int/>
- <sup>76</sup> <https://cites.org/eng>
- <sup>77</sup> <https://www.cms.int/>
- <sup>78</sup> Appendix I taxa, article III paragraph 4 of the convention text, available at: [https://www.cms.int/sites/default/files/instrument/CMS-text.en\\_.PDF](https://www.cms.int/sites/default/files/instrument/CMS-text.en_.PDF)
- <sup>79</sup> <https://accobams.org/>
- <sup>80</sup> <https://www.ascobans.org/>
- <sup>81</sup> <https://www.cms.int/en/document/conservation-plan-migratory-sharks>
- <sup>82</sup> <https://www.cms.int/en/document/single-species-action-plan-angelshark-squatina-squatina-mediterranean-sea>
- <sup>83</sup> Prideaux, G. (2017). Technical Support Information to the CMS Family Guidelines on Environmental Impact Assessments for Marine Noise-generating Activities. Convention on Migratory Species of Wild Animals, Bonn.